

NO SUPERLUMINAL EXPANSION OF THE UNIVERSE*

Don N. Page
CIAR Cosmology Program
Theoretical Physics Institute
Department of Physics
University of Alberta
Edmonton, Alberta
Canada T6G 2J1
Internet: don@page.phys.ualberta.ca

1993 March 2

Abstract

The apparent superluminal growth in the size of a sufficiently large part of the universe can be ascribed to the special relativistic effect of time dilation.

*Alberta Thy-13-93, gr-qc/9303008

In flat (Minkowski) spacetime, special relativity forbids signals and material objects from moving faster than the speed of light c . In any inertial (Lorentz) frame, the linear size (length) of an object cannot expand faster than $2c$, the limit being given by the case in which the two ends of the object each move directly apart at the speed of light. Any faster growth would be superluminal and would not be allowed by special relativity for an object with definite material ends.

(If the material defining the ends can change, there is no such limitation. For example, if a lake freezes over nearly simultaneously, the size of the ice could grow arbitrarily rapidly. As a function of time while the water freezes, the ends of the ice are not definite water molecules but rather growing sequences of molecules which at each instant are at the two ends of the ice. Another example could be the size of a shadow. We shall henceforth exclude such cases from consideration and only consider objects with definite ends that cannot move faster than light as seen locally.)

On the other hand, a sufficiently large region of the universe, with ends that are definite physical entities such as elementary particles or galaxies, apparently can expand much faster than $2c$. For example, a spatially flat homogeneous isotropic cosmological model has the $k = 0$ Friedmann-Robertson-Walker metric

$$ds^2 = -c^2 dt^2 + a^2(t)(dx^2 + dy^2 + dz^2). \quad (1)$$

The particles in this idealized model (which remarkably seems to be an excellent approximation for our universe on length scales of one to ten or so billion light years) each stay at fixed comoving coordinates (x, y, z) as their proper time t increases. Thus the particles are comoving with the spatial coordinates and hence may be considered to be locally at rest with respect to these coordinates.

However, the physical distance between two particles, as measured along a

geodesic of a hypersurface of constant t , is

$$D_t = D_c a(t) \quad (2)$$

where

$$D_c = \sqrt{\Delta x^2 + \Delta y^2 + \Delta z^2} \quad (3)$$

is the (constant) comoving coordinate distance between the two comoving particles, with $(\Delta x, \Delta y, \Delta z)$ being the differences between their comoving coordinates. The physical distance grows at the rate

$$v_t = \frac{d}{dt} D_t = D_c \frac{da}{dt} = H D_t, \quad (4)$$

where

$$H = \frac{1}{a} \frac{da}{dt} \quad (5)$$

is the Hubble expansion rate of the universe at the time t , the logarithmic rate of linear expansion of any comoving part (a part with fixed comoving coordinates x, y, z), with the dimension of reciprocal time.

For the spatially flat model (1) (as also for any open cosmological model), there is no upper limit to the coordinate distance D_c , and hence also to the physical distance D_t , between particles at any fixed t . Thus there is no upper limit to the speed v_t of expansion given by Eq. (4). In particular, it can be much larger than the usual special relativistic limit $2c$. How is this possible?

One idea is that the particles are not actually moving apart, but that space between the particles is expanding by a general relativistic (curved spacetime) effect. This interpretation is certainly natural for the cosmological model (1) written in terms of comoving coordinates, but is it distinct from ordinary expansion in special relativity (flat spacetime)?

To test this, we can use comoving coordinates to examine the expansion of the distance between particles in the flat Minkowski spacetime of special relativity. In inertial coordinates (T, X, Y, Z) , the metric of flat spacetime is

$$ds^2 = -c^2 dT^2 + dX^2 + dY^2 + dZ^2. \quad (6)$$

Suppose we consider a set of particles which fill the region $|X| \leq |T|$ and which have fixed

$$x = \tanh^{-1} \frac{X}{cT} = \frac{1}{2} \ln \frac{cT + X}{cT - X}, \quad (7)$$

$$y = Y, \quad (8)$$

$$z = Z, \quad (9)$$

which serve as comoving coordinates for these particles. At $T = 0$, the particles are all confined to the (Y, Z) plane ($X = 0$), but then each spreads out perpendicularly from there with constant velocity

$$v = \frac{X}{T} = c \tanh x. \quad (10)$$

The proper time along each particle worldline is

$$t = \sqrt{1 - v^2/c^2} T = \sqrt{T^2 - X^2/c^2}. \quad (11)$$

If we invert the relations (7)-(9) and (11), we get the inertial coordinates (T, X, Y, Z) in terms of the comoving coordinates (t, x, y, z) :

$$T = t \cosh x, \quad (12)$$

$$X = ct \sinh x, \quad (13)$$

$$Y = y, \quad (14)$$

$$Z = z. \quad (15)$$

When we differentiate these and insert them into the Minkowski metric (6), we get the metric

$$ds^2 = -c^2 dt^2 + c^2 t^2 dx^2 + dy^2 + dz^2 \quad (16)$$

for the $|X| \leq |T|$ region of the same flat spacetime, but written in the comoving coordinates of the spreading particles.

In these comoving coordinates, the physical distance between two particles with the same y and z , as measured along a geodesic (constant y and z) of a hypersurface of constant t , is (assuming $\Delta x \geq 0$)

$$D_t = ct\Delta x. \quad (17)$$

This grows at the rate

$$v_t = \frac{dD_t}{dt} = c\Delta x. \quad (18)$$

Again, this has no upper limit (e.g., $2c$), since Δx can be arbitrarily large. We seem to have the possibility of an arbitrarily rapid expansion even within the special relativity of flat spacetime! What is going wrong?

The first clue to what is happening is the fact that the hypersurface of constant t (constant proper time from the two-surface $T = 0, X = 0$) is not an (extrinsically) flat hypersurface in the flat spacetime. That is, Eq. (11) shows that the $t = \text{const}$ surface is curved in terms of the inertial coordinates (T, X, Y, Z) . Thus geodesics of that hypersurface are not necessarily geodesics of the enveloping spacetime; the hypersurface $t = \text{const.}$ forces any geodesic within it to bend in the (T, X) plane and therefore not be a geodesic of the full spacetime (unless the geodesic has x , and hence both T and X , constant). This implies that the physical distance D_t within the constant t hypersurface, given by Eq. (17), is not the geodesic distance within the spacetime.

This situation is somewhat analogous to measuring distance on the earth along a great circle, which is a geodesic of the surface (idealized as a sphere), but not of three-dimensional space. In particular, the distance on the surface between the poles is πR , whereas on a geodesic through the space in the middle it is $2R$, where R is the radius of the earth. If the earth were expanding at the speed of light, $dR/dt = c$, the geodesic distance through the center would be growing at the special relativistic limit $2c$, but the great circle distance on the surface would grow at πc , exceeding the limit. Thus we can easily get superluminal expansions if we measure distances that are not geodesics of the whole spacetime.

However, if we go back to our special relativistic model above, we find that the spacetime geodesic distance between the two events at which the two particles (with the same y and z) have the same proper time t is

$$D = \sqrt{\Delta X^2 - c^2 \Delta T^2} = 2ct \sinh \frac{1}{2} \Delta x, \quad (19)$$

which is *greater* than the curved distance D_t given by Eq. (17), increasing the superluminal expansion. How is it possible that the curved path, within the bent $t = \text{const.}$ hypersurface, could be shorter?

In a Riemannian space, with a positive-definite metric (so that all directions are spacelike rather than timelike), the shortest curve between two points is a geodesic, so it cannot be shortened. But in a pseudo-Riemannian space, such as spacetime with its indefinite metric with timelike as well as spacelike directions, any spacelike curve can be shortened by bending it in the timelike direction so that it is more nearly lightlike or null (in the limit of which it would have zero length). The geodesics of the $t = \text{const.}$ hypersurface, which have length D_t , are curved in the timelike direction and hence are shorter than the geodesics of the enveloping spacetime, which have length D . The same would be true in the Friedmann-Robertson-Walker metric (1)

as in the flat metric (16).

Thus our first clue led to a correction (using the length of geodesics of spacetime rather than of a particular spatial hypersurface) that worsens the problem of superluminal expansion. But now that we have a preferred curve (the spacetime geodesic) joining two points on the two particle worldlines, we can ask directly how fast these worldlines are moving apart. If the worldlines are orthogonal to the geodesic joining them, another geodesic joining two slightly later points on the worldlines will have the same length (to first order in the time separation of the two successive joining geodesics), so then the geodesic length between the worldlines is not expanding. This is as true in the curved spacetime of general relativity as in the flat spacetime of special relativity, since it is essentially a consequence of the fact that a geodesic is a curve of extremal length (meaning that with fixed endpoints, its length does not change to first order in a perturbation of the curve, though the extremum need not be a maximum or a minimum and indeed is not for a spacelike geodesic, which gives a saddle point for the distance). Therefore, whether in flat spacetime or in curved spacetime, in no case does the space expand between two worldlines which are orthogonal to geodesics joining them.

If the worldlines are not orthogonal to a succession of geodesics joining corresponding points along them, then the rate of expansion is given by the sum of the outward components of the two worldline velocities (the projections parallel to the geodesic joining the worldlines at the time the expansion is measured), whether in flat spacetime or in curved. Here we finally come to the crucial question, which is whether these velocities and rates are with respect to time T measured by observers moving orthogonally to the joining geodesic, or with respect to the proper time t of the worldlines themselves. It is the former observers that would see the geodesic as existing in one simultaneous moment and whose geodesic length is a purely space-

like distance. The time T of the observers orthogonal to the joining geodesic will be dilated with respect to the proper time t of the particle worldlines by the relativistic “gamma” factor

$$\gamma = \frac{dT}{dt} = (1 - \frac{v^2}{c^2})^{-1/2} \geq 1, \quad (20)$$

where v is the relative speed of the particle worldlines with respect to the orthogonal observers.

It is the derivative of the spacetime geodesic length D with respect to the time T of observers moving orthogonally to that geodesic at its ends that is the sum of the outward components of the ordinary 3-velocities of the ends with respect to the orthogonal observers and hence cannot be larger than twice the speed of light, the luminal limit

$$\frac{dD}{dT} = \mathbf{v}_1 \cdot \mathbf{n}_1 + \mathbf{v}_2 \cdot \mathbf{n}_2 < 2c, \quad (21)$$

where $\mathbf{v}_1$ and $\mathbf{v}_2$ are the ordinary 3-velocities of the ends, and $\mathbf{n}_1$ and $\mathbf{n}_2$ are the unit outward spacelike normal vectors along the joining geodesic. This inequality applies to curved spacetime as well as to flat spacetime. Therefore, if dD/dT is used as the definition of the expansion rate of any region of the universe that has definite physical entities at the ends (so that the ends cannot move faster than light locally), that region, no matter how large, cannot expand superluminally.

On the other hand, the derivative of D with respect to the proper time t of the particle worldlines at the ends of the joining geodesic is the sum of the outward components of the 4-velocities $\mathbf{u}_1$ and $\mathbf{u}_2$ of the ends,

$$\frac{dD}{dt} = \mathbf{u}_1 \cdot \mathbf{n}_1 + \mathbf{u}_2 \cdot \mathbf{n}_2 = \gamma_1 \mathbf{v}_1 \cdot \mathbf{n}_1 + \gamma_2 \mathbf{v}_2 \cdot \mathbf{n}_2, \quad (22)$$

which can exceed $2c$. Similarly, even though the physical distance D_t along a geodesic of the (curved) spacelike hypersurface of fixed t is generally smaller than D , its

derivative with respect to the proper time t can exceed $2c$, in which case one has

$$2c < \frac{dD_t}{dt} < \frac{dD}{dt} < (\gamma_1 + \gamma_2)c. \quad (23)$$

Thus we see that $(\gamma_1 + \gamma_2)c$, rather than $2c$, is the luminal limit when the expansion rate is measured with respect to the proper time t of the ends, as one normally does in calculating cosmological expansion rates, such as v_t given by Eq. (4) above.

In conclusion, we see that no region of the universe, with definite physical entities as ends, can expand faster than $2c$ with respect to the time as measured by observers moving orthogonally to the spacetime geodesics joining the two ends. With respect to the proper time of the ends, the expansion can in principle be as large as $(\gamma_1 + \gamma_2)c$, which is larger than $2c$, but this is simply the time dilation effect of special relativity, just as applicable in the curved spacetime of the universe as in flat spacetime.

This work was supported in part by the Natural Sciences and Engineering Research Council of Canada.